

Dear Members,

LIBTAYO[®] (cemiplimab-rwlc) is now approved for the **adjuvant treatment of adult patients with cutaneous squamous cell carcinoma (CSCC) at high risk of recurrence after surgery and radiation.**

LIBTAYO is also indicated for the treatment of adult patients with metastatic cutaneous squamous cell carcinoma (mCSCC) or locally advanced CSCC (laCSCC) who are not candidates for curative surgery or curative radiation.

Please click [here](#) to learn more.



May help eligible patients access LIBTAYO and navigate the health insurance process. LIBTAYO Surround provides resources, support, and a dedicated team to help patients throughout their treatment journey.

For any questions or concerns, or to report side effects with a Regeneron product for patients enrolled in LIBTAYO Surround, please contact LIBTAYO Surround at **1.877.LIBTAYO** (1.877.542.8296) **Option 1**, Monday–Friday, 8 AM–8 PM Eastern time.

Indications and Usage

LIBTAYO in combination with platinum-based chemotherapy is indicated for the first-line treatment of adult patients with non-small cell lung cancer (NSCLC) with no EGFR, ALK or ROS1 aberrations and is locally advanced where patients are not candidates for surgical resection or definitive chemoradiation OR metastatic.

LIBTAYO as a single agent is indicated for the first-line treatment of adult patients with NSCLC whose tumors have high PD-L1 expression (tumor proportion score [TPS] $\geq 50\%$) as determined by an FDA-approved test, with no EGFR, ALK or ROS1 aberrations, and is locally advanced where patients are not candidates for surgical resection or definitive chemoradiation OR metastatic.

LIBTAYO is indicated for the treatment of adult patients with metastatic cutaneous squamous cell carcinoma (mCSCC) or locally advanced CSCC (laCSCC) who are not candidates for curative surgery or curative radiation.

LIBTAYO is indicated for the adjuvant treatment of adult patients with CSCC at high risk of recurrence after surgery and radiation.

LIBTAYO is indicated for the treatment of adult patients with locally advanced or metastatic basal cell carcinoma (laBCC or mBCC) who have been previously treated with a hedgehog pathway inhibitor or for whom a hedgehog pathway inhibitor is not appropriate.

Important Safety Information

Warnings and Precautions

Severe and Fatal Immune-Mediated Adverse Reactions

Immune-mediated adverse reactions, which may be severe or fatal, can occur in any organ system or tissue at any time after starting treatment. While immune-mediated adverse reactions usually occur

during treatment, they can also occur after discontinuation. Immune-mediated adverse reactions affecting more than one body system can occur simultaneously. Early identification and management are essential to ensuring safe use of PD-1/PD-L1–blocking antibodies. The definition of immune-mediated adverse reactions included the required use of systemic corticosteroids or other immunosuppressants and the absence of a clear alternate etiology. Monitor closely for symptoms and signs that may be clinical manifestations of underlying immune-mediated adverse reactions. Evaluate liver enzymes, creatinine, and thyroid function at baseline and periodically during treatment. In cases of suspected immune-mediated adverse reactions, initiate appropriate workup to exclude alternative etiologies, including infection. Institute medical management promptly, including specialty consultation as appropriate.

Withhold or permanently discontinue LIBTAYO depending on severity of the adverse reaction (see Section 2 Dosage and Administration in the accompanying Full Prescribing Information). In general, if LIBTAYO requires interruption or discontinuation, administer systemic corticosteroid therapy (1 to 2 mg/kg/day prednisone or equivalent) until improvement to Grade 1 or less. Upon improvement to Grade 1 or less, initiate corticosteroid taper and continue to taper over at least 1 month. Consider administration of other systemic immunosuppressants in patients whose immune-mediated adverse reactions are not controlled with corticosteroids. Toxicity management guidelines for adverse reactions that do not necessarily require systemic steroids (e.g., endocrinopathies and dermatologic reactions) are discussed below.

The incidence and severity of immune-mediated adverse reactions were similar when LIBTAYO was administered as a single agent or in combination with chemotherapy.

Immune-mediated pneumonitis: LIBTAYO can cause immune-mediated pneumonitis. In patients treated with other PD-1/PD-L1–blocking antibodies, the incidence of pneumonitis is higher in patients who have received prior thoracic radiation. Immune-mediated pneumonitis occurred in 2.6% (33/1281) of patients receiving LIBTAYO, including Grade 4 (0.3%), Grade 3 (0.6%), and Grade 2 (1.6%).

Pneumonitis led to permanent discontinuation in 1.3% of patients and withholding of LIBTAYO in 1.4% of patients. Systemic corticosteroids were required in all patients with pneumonitis. Pneumonitis resolved in 61% of the 33 patients. Of the 18 patients in whom LIBTAYO was withheld, 10 reinitiated after symptom improvement; of these, 4/10 (40%) had recurrence of pneumonitis.

Immune-mediated colitis: LIBTAYO can cause immune-mediated colitis. The primary component of immune-mediated colitis was diarrhea. Cytomegalovirus (CMV) infection/reactivation has been reported in patients with corticosteroid-refractory immune-mediated colitis treated with PD-1/PD-L1–blocking antibodies. In cases of corticosteroid-refractory immune-mediated colitis, consider repeating infectious workup to exclude alternative etiologies. Immune-mediated colitis occurred in 2% (25/1281) of patients receiving LIBTAYO, including Grade 3 (0.8%) and Grade 2 (0.9%). Colitis led to permanent discontinuation in 0.4% of patients and withholding of LIBTAYO in 1.2% of patients. Systemic corticosteroids were required in all patients with colitis. Colitis resolved in 56% of the 25 patients. Of the 16 patients in whom LIBTAYO was withheld, 6 reinitiated LIBTAYO after symptom improvement; of these, 4/6 (67%) had recurrence.

Immune-mediated hepatitis: LIBTAYO can cause immune-mediated hepatitis. Immune-mediated hepatitis occurred in 2.4% (31/1281) of patients receiving LIBTAYO, including fatal (<0.1%), Grade 4 (0.3%), Grade 3 (1.6%), and Grade 2 (0.2%). Hepatitis led to permanent discontinuation of LIBTAYO in 1.4% of patients and withholding of LIBTAYO in 0.7% of patients. Systemic corticosteroids were required in all patients with hepatitis. Additional immunosuppression with mycophenolate was required in 13% (4/31) of these patients. Hepatitis resolved in 39% of the 31 patients. Of the 9 patients in whom LIBTAYO was withheld, 5 reinitiated LIBTAYO after symptom improvement; of these, 1/5 (20%) had recurrence.

Immune-mediated endocrinopathies:

- **Adrenal insufficiency:** LIBTAYO can cause primary or secondary adrenal insufficiency. For Grade 2 or higher adrenal insufficiency, initiate symptomatic treatment, including hormone replacement as clinically indicated. Withhold LIBTAYO depending on severity. Adrenal insufficiency occurred in 0.5% (6/1281) of patients receiving LIBTAYO, including Grade 3 (0.5%). Adrenal insufficiency led to permanent discontinuation of LIBTAYO in 1 (<0.1%) patient. LIBTAYO was withheld in 1 (<0.1%) patient due to adrenal insufficiency and not reinitiated. Systemic

corticosteroids were required in 83% (5/6) patients with adrenal insufficiency; of these, the majority remained on systemic corticosteroids. Adrenal insufficiency had resolved in 17% of the 6 patients

- **Hypophysitis:** LIBTAYO can cause immune-mediated hypophysitis. Hypophysitis can present with acute symptoms associated with mass effect such as headache, photophobia, or visual field defects. Hypophysitis can cause hypopituitarism. Initiate hormone replacement as clinically indicated. Withhold or permanently discontinue depending on severity. Hypophysitis occurred in 0.5% (7/1281) of patients receiving LIBTAYO, including Grade 3 (0.2%) and Grade 2 (0.3%) adverse reactions. Hypophysitis led to permanent discontinuation of LIBTAYO in 1 (<0.1%) patient and withholding of LIBTAYO in 2 (0.2%) patients. Systemic corticosteroids were required in 86% (6/7) of patients with hypophysitis. Hypophysitis resolved in 14% of the 7 patients. Of the 2 patients in whom LIBTAYO was withheld for hypophysitis, none of the patients reinitiated
- **Thyroid disorders:** LIBTAYO can cause immune-mediated thyroid disorders. Thyroiditis can present with or without endocrinopathy. Hypothyroidism can follow hyperthyroidism. Initiate hormone replacement or medical management as clinically indicated. Withhold or permanently discontinue LIBTAYO depending on severity
- **Thyroiditis:** Thyroiditis occurred in 0.6% (8/1281) of patients receiving LIBTAYO, including Grade 2 (0.3%) adverse reactions. No patient discontinued LIBTAYO due to thyroiditis. Thyroiditis led to withholding of LIBTAYO in 1 (<0.1%) patient. Systemic corticosteroids were not required in any patient with thyroiditis. Thyroiditis resolved in 13% of the 8 patients. Blood thyroid stimulating hormone increased and blood thyroid stimulating hormone decreased have also been reported
- **Hyperthyroidism:** Hyperthyroidism occurred in 3% (39/1281) of patients receiving LIBTAYO, including Grade 3 (<0.1%) and Grade 2 (0.9%). No patient discontinued treatment and LIBTAYO was withheld in 7 (0.5%) patients due to hyperthyroidism. Systemic corticosteroids were required in 8% (3/39) of patients. Hyperthyroidism resolved in 56% of 39 patients. Of the 7 patients in whom LIBTAYO was withheld for hyperthyroidism, 2 patients reinitiated LIBTAYO after symptom improvement; of these, none had recurrence of hyperthyroidism
- **Hypothyroidism:** Hypothyroidism occurred in 7% (87/1281) of patients receiving LIBTAYO, including Grade 3 (<0.1%) and Grade 2 (6%). Hypothyroidism led to permanent discontinuation of LIBTAYO in 3 (0.2%) patients. Hypothyroidism led to withholding of LIBTAYO in 9 (0.7%) patients. Systemic corticosteroids were required in 1.1% (1/87) of patients with hypothyroidism. Hypothyroidism resolved in 6% of the 87 patients. Majority of the patients with hypothyroidism required long-term thyroid hormone replacement. Of the 9 patients in whom LIBTAYO was withheld for hypothyroidism, 1 reinitiated LIBTAYO after symptom improvement and did not have recurrence of hypothyroidism
- **Type 1 diabetes mellitus, which can present with diabetic ketoacidosis:** Monitor for hyperglycemia or other signs and symptoms of diabetes. Initiate treatment with insulin as clinically indicated. Withhold LIBTAYO depending on severity. Type 1 diabetes mellitus occurred in <0.1% (1/1281) of patients (Grade 4). No patient discontinued treatment due to Type 1 diabetes mellitus. Type 1 diabetes mellitus led to withholding of LIBTAYO in 0.1% of patients, treatment was reinitiated after symptom improvement. Patient received long-term insulin therapy

Immune-mediated nephritis with renal dysfunction: LIBTAYO can cause immune-mediated nephritis. Immune-mediated nephritis occurred in 0.7% (9/1281) of patients receiving LIBTAYO, including fatal (<0.1%), Grade 3 (<0.1%), and Grade 2 (0.5%). Nephritis led to permanent discontinuation in 0.2% of patients and withholding of LIBTAYO in 0.4% of patients. Systemic corticosteroids were required in all patients with nephritis. Nephritis resolved in 78% of the 9 patients. Of the 5 patients in whom LIBTAYO was withheld, 4 reinitiated LIBTAYO after symptom improvement; of these, 1/4 (25%) had recurrence.

Immune-mediated dermatologic adverse reactions: LIBTAYO can cause immune-mediated rash or dermatitis. Exfoliative dermatitis, including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug rash with eosinophilia and systemic symptoms (DRESS) has occurred with PD-1/PD-L1–blocking antibodies. Immune-mediated dermatologic adverse reactions occurred in 1.9% (24/1281) of patients receiving LIBTAYO, including Grade 3 (0.9%) and Grade 2 (0.8%). Immune-mediated dermatologic adverse reactions led to permanent discontinuation in 0.2% of patients and withholding of LIBTAYO in 1.3% of patients. Systemic corticosteroids were required in all patients with immune-

mediated dermatologic adverse reactions. Immune-mediated dermatologic adverse reactions resolved in 71% of the 24 patients. Of the 17 patients in whom LIBTAYO was withheld for dermatologic adverse reaction, 13 reinitiated LIBTAYO after symptom improvement; of these, 5/13 (38%) had recurrence of the dermatologic adverse reaction. Topical emollients and/or topical corticosteroids may be adequate to treat mild to moderate non-exfoliative rashes.

Other immune-mediated adverse reactions: The following clinically significant immune-mediated adverse reactions occurred at an incidence of <1% in 1281 patients who received LIBTAYO or were reported with the use of other PD-1/PD-L1–blocking antibodies. Severe or fatal cases have been reported for some of these adverse reactions.

- **Cardiac/vascular:** Myocarditis, pericarditis, and vasculitis. Permanently discontinue for Grades 2, 3, or 4 myocarditis
- **Nervous system:** Meningitis, encephalitis, myelitis and demyelination, myasthenic syndrome/myasthenia gravis (including exacerbation), Guillain-Barré syndrome, nerve paresis, and autoimmune neuropathy
- **Ocular:** Uveitis, iritis, and other ocular inflammatory toxicities. Some cases can be associated with retinal detachment. Various grades of visual impairment to include blindness can occur. If uveitis occurs in combination with other immune-mediated adverse reactions, consider a Vogt-Koyanagi-Harada–like syndrome, as this may require treatment with systemic steroids to reduce the risk of permanent vision loss
- **Gastrointestinal:** Pancreatitis to include increases in serum amylase and lipase levels, gastritis, duodenitis, stomatitis
- **Musculoskeletal and connective tissue:** Myositis/polymyositis/dermatomyositis, rhabdomyolysis, and associated sequelae including renal failure, arthritis, polymyalgia rheumatica
- **Endocrine:** Hypoparathyroidism
- **Other (hematologic/immune):** Hemolytic anemia, aplastic anemia, hemophagocytic lymphohistiocytosis (HLH), systemic inflammatory response syndrome, histiocytic necrotizing lymphadenitis (Kikuchi lymphadenitis), sarcoidosis, immune thrombocytopenia, solid organ transplant rejection, other transplant (including corneal graft) rejection

Infusion-Related Reactions

Severe or life-threatening infusion-related reactions occurred in 0.2% of patients receiving LIBTAYO as a single agent. Monitor patients for signs and symptoms of infusion-related reactions. Common symptoms of infusion-related reaction include nausea, pyrexia, and vomiting. Interrupt or slow the rate of infusion or permanently discontinue LIBTAYO based on severity of reaction.

Complications of Allogeneic HSCT

Fatal and other serious complications can occur in patients who receive allogeneic hematopoietic stem cell transplantation (HSCT) before or after being treated with a PD-1/PD-L1–blocking antibody. Transplant-related complications include hyperacute graft-versus-host disease (GVHD), acute GVHD, chronic GVHD, hepatic veno-occlusive disease (VOD) after reduced intensity conditioning, and steroid-requiring febrile syndrome (without an identified infectious cause). These complications may occur despite intervening therapy between PD-1/PD-L1 blockade and allogeneic HSCT. Follow patients closely for evidence of transplant-related complications and intervene promptly. Consider the benefit versus risks of treatment with a PD-1/PD-L1–blocking antibody prior to or after an allogeneic HSCT.

Embryo-Fetal Toxicity

LIBTAYO can cause fetal harm when administered to a pregnant woman due to an increased risk of immune-mediated rejection of the developing fetus resulting in fetal death. Advise women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with LIBTAYO and for at least 4 months after the last dose.

Adverse Reactions

The most common adverse reactions (≥15%) are:

- *As a single agent in mCSCC or laCSCC, mBCC or laBCC, and NSCLC with high PD-L1 expression:* fatigue, musculoskeletal pain, rash, diarrhea, and anemia
- *As a single agent in adjuvant CSCC at high risk of recurrence:* rash and pruritus

- *In combination with platinum-based chemotherapy in NSCLC:* alopecia, musculoskeletal pain, nausea, fatigue, peripheral neuropathy, and decreased appetite

Use in Specific Populations

- **Lactation:** Because of the potential for serious adverse reactions in breastfed children, advise women not to breastfeed during treatment and for at least 4 months after the last dose of LIBTAYO
- **Females and males of reproductive potential:** Verify pregnancy status in females of reproductive potential prior to initiating LIBTAYO

Please click [here](#) for full Prescribing Information.

For product information about LIBTAYO, click [here](#). For patient support resources, visit [LIBTAYO Surround](#).

Thank you,



Melissa Souders, PACS

Sr. Reimbursement Business Manager

South Central

C: 214-564-3024

W: 469-516-8961

E: Melissa.Souders@regeneron.com

REGENERON
SCIENCE TO MEDICINE®

For Colorado Prescribers – [Click Here for Pricing Information](#)

For Connecticut Prescribers – [Click Here for Pricing Information](#)

If you no longer wish to receive LIBTAYO emails, please reply to this message with "Unsubscribe" in the subject line.

Reference: LIBTAYO (cemiplimab-rwlc) injection full U.S. Prescribing Information. Regeneron Pharmaceuticals, Inc.

If you have a question about a Regeneron product, please call 1-844-643-7346.

© 2025 Regeneron Pharmaceuticals, Inc. All rights reserved.

777 Old Saw Mill River Rd, Tarrytown, NY 10591

US.LIB.25.05.0013 10/25

Regeneron - Internal

This e-mail and any attachment hereto, is intended only for use by the addressee(s) named above and may contain legally privileged and/or confidential information. If you are not the intended recipient of this e-mail, any dissemination, distribution or copying of this email, or any attachment hereto, is strictly prohibited. If you receive this email in error please immediately notify me by return electronic mail and permanently delete this email and any attachment hereto, any copy of this e-mail and of any such attachment, and any printout thereof. Finally, please note that only authorized representatives of Regeneron Pharmaceuticals, Inc. have the power and authority to enter into business dealings with any third party.
